

R. JOHN ELLIS, JR., M.D.
LAWRENCE A. SCHAPER, M.D.
MARK G. SMITH, M.D.
G. JEFFREY POPHAM, M.D.
AKBAR NAWAB, M.D.
MICHAEL L. SALAMON, M.D.
MATTHEW R. PRICE, M.D.



DANIEL E. RUEFF, M.D.
SEAN M. GRIFFIN, M.D.
BRENT J. SINICROPE, M.D.
EVAN RHEA, M.D.
KELSI J. BARNES, PA-C
JENNIFER R. STREET, PA-C

(MACI) MATRIX-INDUCED AUTOLOGOUS CHONDROCYTE IMPLANT FEMORAL CONDYLE PROTOCOL (Dr. Akbar Nawab)

GENERAL GUIDELINES: (0-3 Months following surgery)

- Patients typically start formal PT at one week after surgery; patients will work towards pain-free and full passive knee extension with limited weight-bearing. Safety with crutches/walker will need to be assessed pre and post-op. Over time, the goal is to be free of ambulation devices and knee braces while becoming thoroughly independent with rehabilitation exercises.

Week 1

- CPM 0-30° (12-24 hours after surgery) and utilize at least one hour each day. (if issued)
- **ROM 0-90° (passively)**
- Check incision and swelling assessment. Check for DVT.
- Use crutches at all times and <20% weight bearing
- Brace locked in full extension with sleeping and walking. Unlocked 0-30° with transfers sitting and NWB ADL's.
- Elevate the entire leg. Do not place pillows under the knee for long periods of time.
- Restore knee extension (low load-long duration stretching).
- Hamstring/gastrocnemius stretches.
- Ankle ROM for swelling and DVT prevention.
- ROM of knee (assisted heel slides): 0°-30°
- NMES (with BFR if indicated) for Quadriceps activation and re-education.
- Initiate SLR's and Hip ABD as tolerated with knee extended and no brace.

Weeks 2-3

- Progress to 30% weight bearing with appropriate assistive device (encourage heel-toe gait)
- **ROM 0-90° by end of week 2; ROM to tolerance after week 3**
- Restore patellar mobility as incision heals
- Initiate remainder of open chain straight leg exercises. *Focus on the knee remaining locked in the concentric and eccentric phase. Increase reps and hold time initially, before adding resistance.*
- Restore knee extension (low load-long duration stretching).
- Continue with NMES, BFR, open chain strengthening/stretching and other modalities for swelling and DVT prevention per PT POC

EASTPOINT OFFICE
13151 MAGISTERIAL DRIVE, SUITE 200
LOUISVILLE, KENTUCKY 40223
Telephone 502-587-1236
Fax 502-587-0126

BOOKKEEPING DEPARTMENT
5120 DIXIE HIGHWAY, SUITE 103
LOUISVILLE, KENTUCKY 40216
Telephone 502-587-7269
Fax 502-587-0318

SOUTHEND OFFICE
5120 DIXIE HIGHWAY, SUITE 103
LOUISVILLE, KENTUCKY 40216
Telephone 502-449-0449
Fax 502-449-3277

Weeks 4-6

- Progress to 40%-60% weight bearing with 1-2 crutches.
- **ROM to tolerance**
- May unlock brace 0°-60° at week 4 and increase to full flexion by week 6
- Normalize gait with full knee ROM in the brace by week 6
- Initiate cycle for ROM as tolerated with PF protection
- Focus on an independent home exercise program.
- Perform simple ADLs with assistance of crutches (navigating stairs, showering, etc.)

Weeks 7-12

- Initially progress weight bearing to 80% at week 7 and then progress to FWB at 8-10 weeks.
- Go without crutches indoors and one crutch if outdoors (or unfamiliar areas)
- Discontinue brace at week 7 if no extension lag with SLR.
- Regain full ROM
- Continue stationary cycle
- Quadriceps stretching weeks 9-10
- Initiate closed chain exercises; avoid PF pain to focus on strength and proprioception.
- Start driving again and return to office or seated work.

3-6 Months

- Increase resistance with all exercises as tolerated.
- Full and pain-free AROM, no brace, and no assistive devices.
- Increase leg strength to allow for normal gait and walking longer distances.
- Leg press 0-60° and progress to 0-90°
- Forward lunges and wall squats
- Return to low impact recreational activities including: cycling, yoga & Pilates, swimming, elliptical/treadmill, golf, rowing/kayaking, and dancing.
- Return to a more physically active job such as nursing or construction.
- Return to daily activities that require strength/endurance.

6-9 Months

- Agility and balance drills can be initiated.
- Walking >3 miles
- Continue with swimming, cycling, and elliptical and increase distance for endurance as tolerated

9-12 Months

- Jogging, running, and high impact aerobics are allowed
- High impact sports such as tennis, football, basketball, skiing and snowboarding are allowed at 12+ months